

Kleinian groups with discrete length spectrum

Richard D. Canary*
Christopher J. Leininger†

February 2, 2008

Given a Kleinian group $\Gamma \subset \mathrm{PSL}_2(\mathbb{C}) \cong \mathrm{Isom}_+(\mathbb{H}^3)$, we define the *real length spectrum of Γ without multiplicities* to be the set of all real translation lengths of elements of Γ acting on $\mathbb{H}^3$

$$\mathfrak{L}(\Gamma) = \{l(x) \mid x \in \Gamma\} \subset \mathbb{R}$$

Equivalently, these are the lengths of closed geodesics in the quotient orbifold.

In this paper we characterize finitely generated, torsion-free Kleinian groups with discrete real length spectrum. In the process, we obtain a sharper version of the Covering Theorem which may be of independent interest.

Theorem 1 *Suppose that Γ is a finitely generated, torsion-free Kleinian group. Then the real length spectrum $\mathfrak{L}(\Gamma)$ is discrete if and only if either*

1. Γ is geometrically finite,
2. there exists a hyperbolic 3-manifold $M = \mathbb{H}^3/G$ which fibers over the circle and Γ is a fiber subgroup of G , or
3. there exists a hyperbolic 3-manifold $M = \mathbb{H}^3/G$ which fibers over $S^1/\langle z \mapsto \bar{z} \rangle$ and Γ is a singular fiber subgroup of G .

We recall that if a hyperbolic 3-manifold $M = \mathbb{H}^3/G$ fibers over a 1-orbifold Q , then a fiber subgroup of G is the fundamental group of the pre-image of a regular point of Q , while a singular fiber subgroup is the fundamental group of the pre-image of a singular point of Q . (Notice that one only really obtains a conjugacy class of subgroups.)

Proof: We refer the reader to [Ca1] and [Ca2] for details and terminology. Let $N = \mathbb{H}^3/\Gamma$, let $\epsilon > 0$ be less than the 3-dimensional Margulis constant, and let N_ϵ^0 denote the complement of the ϵ -cuspidal thin part of N . By the Tameness Theorem (see Agol [Ag] or Calegari-Gabai [CG]), N is topologically tame.

It is well-known that a geometrically finite Kleinian group has discrete real length spectrum. We sketch a brief proof for the reader's convenience. If Γ is

*Research supported in part by NSF grants 0203698 and 0504791

†Research supported in part by an N.S.F. postdoctoral fellowship

geometrically finite, then there exists a compact subset K of $N = \mathbb{H}^3/\Gamma$ such that any closed geodesic intersects K . (One may, for example, take K to be the intersection of the convex core of N with N_ϵ^0 .) If $\mathfrak{L}(\Gamma)$ is indiscrete, then there exists an infinite collection of distinct closed geodesics of length at most L (for some $L > 0$) all of which intersect K . If this were the case, some sequence of closed geodesics in N would have to accumulate at a closed geodesic, which is impossible. Therefore, $\mathfrak{L}(\Gamma)$ is discrete.

Since a fiber subgroup, as described in (2), or singular fiber subgroup, as described in (3), is a subgroup of a geometrically finite Kleinian group, they must also have discrete real length spectrum.

We now assume that Γ is a geometrically infinite Kleinian group with discrete real length spectrum and prove that Γ is either a fiber subgroup or a singular fiber subgroup of the fundamental group of a finite volume hyperbolic 3-manifold which fibers over a 1-orbifold. Let $U \cong \bar{S} \times [0, \infty) \subset N_\epsilon^0$ be a neighborhood of a geometrically infinite end of N_ϵ^0 and let $U_p \cong S \times [0, \infty) \subset N$ be its parabolic extension. By the main result of [Ca1], there exists a sequence of useful simplicial hyperbolic surfaces $f_n : X_n \rightarrow U_p$, each properly homotopic in U_p to $S \times \{0\}$, which exits the end.

As the length spectrum of Γ is discrete, there is a uniform positive lower bound to the length of any closed geodesic in N . Therefore, there is a uniform positive lower bound on the length of any closed geodesic on X_n whose image under f_n is essential and not an accidental parabolic (i.e. not homotopic into a cusp of N). On the other hand, Lemma 7.1 in [Ca1] guarantees that given any $A > 0$, there exists $n_A > 0$ so that for all $n \geq n_A$, any closed geodesics in X_n which is inessential or accidentally parabolic under f_n must have length at least A on X_n . Therefore, there is a uniform lower bound to the length of a closed geodesic on any X_n .

Lemma 7.1 in [Ca2] then gives that there exists $C > 0$ and, for all n , a point x_n in the ϵ_0 -thick part of X_n and a minimal generating set $\{a_1^n, \dots, a_m^n\}$ for $\pi_1(X_n, x_n)$ all of which are represented by curves (based at x_n) of length at most C . By passing to a subsequence, we may assume that all the minimal generating sets are topologically equivalent. Therefore, there exist markings $g_n : S \rightarrow X_n$ so that any chosen finite set of curves on S have representatives with uniformly bounded length on X_n . Composing, we obtain a sequence of representations

$$\rho_n = (f_n \circ g_n)_* : \pi_1(S) \rightarrow \Gamma \subset \mathrm{PSL}_2(\mathbb{C})$$

A result of Kim [Ki] gives that there exists a finite set $\{s_1, \dots, s_k\}$ of elements of $\pi_1(S)$ such that an irreducible representation of $\pi_1(S)$ is determined up to conjugacy by their lengths. Since the lengths of $s_1, \dots, s_k$ with respect to $g_n : S \rightarrow X_n$ are bounded, so are the set of lengths $\rho_n(s_i)$. Since $\mathfrak{L}(\Gamma)$ is discrete, we may pass to a subsequence, again called $\{\rho_n\}$, so that $l(\rho_n(s_i))$ is constant for all i . Therefore, all representations in this sequence are conjugate to ρ_1 . Let $h_n \in \mathrm{PSL}_2(\mathbb{C})$ be the conjugating element, so that $\rho_n = h_n \rho_1 h_n^{-1}$.

We now observe that U_p is incompressible (i.e. that ρ_1 is injective). First note that, since all the ρ_n are conjugate, if $s \in \mathrm{Ker}(\rho_1)$ is non-trivial, then

$s \in \text{Ker}(\rho_n)$ for all n , and that the length of $g_n(s)$ in X_n is uniformly bounded. However, since $f_n(g_n(s))$ is non-essential, $\{l_{X_n}(g_n(s))\}$ converges to ∞ , which is a contradiction.

Finally, consider the cover $\tilde{N}_U$ of N corresponding to $\pi_1(U)$. The conjugating elements h_n descend to isometries $\hat{h}_n$ of $\tilde{N}_U$. Let $\tilde{f}_n : X_n \rightarrow \tilde{N}_U$ denote the lift of f_n to $\tilde{N}_U$ and let $\tilde{x}_n = \tilde{f}_n(x_n)$. Note that there exists $D > 0$ so that if $x \in \tilde{N}_U$ and there exist representatives of $\{\tilde{f}_1(a_1^1), \dots, \tilde{f}_1(a_m^1)\}$ based at x of length at most C , then $d(x, \tilde{x}_1) \leq D$. It then follows that $d(\tilde{x}_n, \hat{h}_n(\tilde{x}_1)) \leq D$ for all n . Since $\{f_n(X_n)\}$ exits U , one sees that $\hat{h}_n(U) \cap U$ is non-empty for all large enough n , and that there are infinitely many distinct $\hat{h}_n$.

Let N_1 be the quotient of $\tilde{N}_U$ by its full group of orientation-preserving isometries. We have established that the covering map $\pi_U : \tilde{N}_U \rightarrow N_1$ is infinite-to-one on U .

We now recall the Covering Theorem (see [T, Theorem 9.2.2], [Ca2] and Agol [Ag]):

Covering Theorem: *Suppose that $\widehat{M}$ is a hyperbolic 3-manifold with finitely generated fundamental group and M is a hyperbolic 3-orbifold. If $p : \widehat{M} \rightarrow M$ is an orbifold cover which is infinite-to-one on a neighborhood U of a geometrically infinite end of $\widehat{M}_\epsilon^0$, then M has finite volume and has a finite manifold cover M' which fibers over the circle such that either*

- a) $\widehat{M}$ is the cover of M' associated to the fiber, or*
- b) $\widehat{M}_0^\epsilon$ is a twisted $\mathbf{R}$ -bundle and $\widehat{M}$ is double covered by the manifold $\widehat{M}'$ which is the cover of M' associated to the fiber.*

The Covering Theorem immediately implies that Γ_U is the fiber group associated to a 3-manifold which fibers over the circle. If $\Gamma = \Gamma_U$, we're done. If not, we apply the Covering Theorem again to the map $p_U : \tilde{N}_U \rightarrow N$ to conclude that p_U is finite-to-one. Hempel's Finite Index Theorem [He, Theorem 10.5] implies that N_0^ϵ is a twisted $\mathbf{R}$ -bundle and that the cover p_U is two-to-one.

The following addendum to the Covering Theorem allows us to conclude that Γ is a singular fiber subgroup, in the case that N_ϵ^0 is a twisted $\mathbf{R}$ -bundle.

Proposition 1 *Suppose that $\widehat{M}$ is a geometrically infinite hyperbolic 3-manifold such that $\widehat{M}_0^\epsilon$ is a twisted $\mathbf{R}$ -bundle over a compact surface, and let $p' : \widehat{M}' \rightarrow \widehat{M}$ be a two-fold cover such that $\widehat{M}'_0^\epsilon$ is an untwisted $\mathbf{R}$ -bundle.*

1) If there is a finite volume hyperbolic 3-manifold M'' that fibers over the circle with covering $q' : \widehat{M}' \rightarrow M''$ corresponding to the fiber, then $\widehat{M}$ covers a finite volume hyperbolic orbifold M .

2) If $p : \widehat{M} \rightarrow M$ is a cover of a finite volume hyperbolic orbifold M , then M has a finite manifold cover $M' = \mathbb{H}^3/G'$ which fibers over the orbifold $S^1/\langle z \mapsto \bar{z} \rangle$ and $\widehat{M}$ is the cover associated to a singular fiber subgroup of G' .

We now turn to the proof of Proposition 1: For the first statement, we note that the covers $p' : \widehat{M}' \rightarrow \widehat{M}$ and $q' : \widehat{M}' \rightarrow M''$ are regular covers with covering groups generated by an involution $a \in \text{Isom}_+(\widehat{M}')$ and infinite order isometry $b \in \text{Isom}_+(\widehat{M}')$, respectively. Since $\text{Isom}_+(\widehat{M}')$ acts properly discontinuously on $\widehat{M}'$ and $\langle b \rangle$ has quotient M'' of finite volume, so the covering $\hat{p} : \widehat{M}' \rightarrow M = \widehat{M}' / \text{Isom}_+(\widehat{M}')$ is onto the finite volume orbifold M . Since $\widehat{M}$ is the quotient of $\widehat{M}'$ by the subgroup $\langle a \rangle < \text{Isom}_+(\widehat{M}')$, it follows that there is a covering $p : \widehat{M} \rightarrow M$ so that $\hat{p} = p \circ p'$, verifying the first statement.

For the second assertion, define $\hat{p} = p \circ p'$. The Covering Theorem, applied to $\hat{p} : \widehat{M}' \rightarrow M$, implies that M has a finite manifold cover M'' which fibers over the circle so that $\widehat{M}'$ is the cover associated to the fiber. Moreover, examining the proof of the Covering Theorem, we see that if $q : M'' \rightarrow M$ and $q' : \widehat{M}' \rightarrow M''$ are the associated covering maps, we may assume that $q \circ q' = \hat{p} = p \circ p'$.

As above, the covers $p' : \widehat{M}' \rightarrow \widehat{M}$ and $q' : \widehat{M}' \rightarrow M''$ are regular with covering group of order two, generated by $a \in \text{Isom}_+(\widehat{M}')$, and infinite cyclic covering group, generated by $b \in \text{Isom}_+(\widehat{M}')$, respectively. Since $\langle b \rangle$ has finite volume quotient M'' and $\text{Isom}_+(\widehat{M}')$ acts properly discontinuously, it follows $\langle b \rangle$ has finite index in $\text{Isom}_+(\widehat{M}')$ and hence in $\langle a, b \rangle$. Therefore, there exists $k > 0$ such that $\langle b^k \rangle$ is normal in $\langle a, b \rangle$.

Since $\langle b^k \rangle$ is normal in $\langle a, b \rangle$, we see that either $ab^ka = b^k$ or $ab^ka = b^{-k}$. We now see that it must be the case that $ab^ka = b^{-k}$. Let T be a connected separating surface in $(\widehat{M}')_\epsilon^0$ which is preserved by a , i.e. $a(T) = T$. If $n > 0$ is chosen large enough then $b^{nk}(T)$ does not intersect T . Since a interchanges the two components of $(N_U)_\epsilon^0 - T$, we see immediately that $ab^{nk}a(T)$ and $b^{nk}(T)$ lie on opposite sides of T . Therefore, $ab^{nk}a \neq b^{nk}$, which implies that $ab^ka \neq b^k$. It follows that $ab^ka = b^{-k}$ and that $\langle a, b^k \rangle \cong \mathbb{Z}_2 * \mathbb{Z}_2$ is generated by a and ab^k .

If n is even, then ab^{nk} is conjugate in $\langle a, b \rangle$ to a . Since a acts freely, so does ab^{nk} . Therefore, $\langle a, b^{2k} \rangle \cong \mathbb{Z}_2 * \mathbb{Z}_2$ acts freely on M' . Let M' be the quotient of $\widehat{M}'$ by the group $\langle a, b^{2k} \rangle$. The main result of Hempel-Jaco [HJ] then implies that M' fibers over the orbifold $S^1 / \langle z \mapsto \bar{z} \rangle$ and that $\widehat{M}'$ is the cover associated to a singular fiber subgroup. (Alternatively, one may assume that k is large enough that $b^k(T)$ does not intersect T and prove directly that the region between T and $b^k(T)$ is a fundamental region for the group $\langle a, b^{2k} \rangle$. One may then fairly explicitly check that the quotient $\widehat{M}' / \langle a, b^{2k} \rangle$ fibers over the orbifold $S^1 / \langle z \mapsto \bar{z} \rangle$ and that $\widehat{M}'$ is a cover associated to a singular fiber subgroup.) One may then check that $\hat{p} : \widehat{M}' \rightarrow M$ descends to a covering map $\hat{p}' : M' \rightarrow M$. This completes the proof of Proposition 1, which in turn completes the proof of our Main Theorem.

If we combine Proposition 1 with our earlier statement of the Covering Theorem, we get the following slightly sharper version.

Covering Theorem(sharper version): Suppose that $\widehat{M}$ is a hyperbolic 3-manifold with finitely generated fundamental group and M is a hyperbolic 3-orbifold. If $p : \widehat{M} \rightarrow M$ is an orbifold cover which is infinite-to-one on a neighborhood U of a geometrically infinite end of $\widehat{M}_\epsilon^0$, then M has finite volume and has a finite manifold cover $M' = \mathbb{H}^3/G'$ such that either

1. M' fibers over the circle and $\widehat{M}$ is the cover associated to a fiber subgroup of G' , or
2. M' fibers over the orbifold $S^1/\langle z \mapsto \bar{z} \rangle$ and $\widehat{M}$ is the cover of M' associated to a singular fiber subgroup of G' .

References

- [Ag] I. Agol, *Tameness of hyperbolic 3-manifolds*, e-print <http://arxiv.org/abs/math.GT/0405568>.
- [CG] D. Calegari and D. Gabai, *Shrinkwrapping and the taming of hyperbolic 3-manifolds*, e-print, <http://arxiv.org/abs/math.GT/0407161>.
- [Ca1] R. D. Canary, *Ends of hyperbolic 3-manifolds*, J. AMS **6**, (1993), 1–35.
- [Ca2] R. D. Canary, *A covering theorem for hyperbolic 3-manifolds and its applications*, Topology **35**, (1996) 751–778.
- [He] J. Hempel, *3-manifolds*, Princeton University Press, 1976.
- [HJ] J. Hempel and W. Jaco, *Fundamental groups of 3-manifolds which are extensions*, Ann. of Math. **95**(1972), 86–98.
- [Ki] I. Kim, *Marked length spectrum on the finite set of elements determines the irreducible representation in the isometry group of rank one symmetric space of noncompact type*, Monogr. Geom. Topology 25, Internat. Press, Cambridge, MA, 1998, 99–107.
- [T] W. P. Thurston, *The Geometry and Topology of Three-Manifolds*, Princeton University course notes, available at <http://www.msri.org/publications/books/gt3m/> (1980).